

## REPORT DOCUMENTATION PAGE

AD-A231 379

1a. REPORT SECURITY CLASSIFICATION (U) 1b. RESTRICTED

2a. SECURITY CLASSIFICATION AUTHORITY NA 2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA

4. PERFORMING ORGANIZATION REPORT NUMBER(S) 5. MONITORING ORGANIZATION REPORT NUMBER(S)

New England Medical Center Hospitals, Inc.

NA

6a. NAME OF PERFORMING ORGANIZATION  
New England Medical Center  
Hospitals, Inc.6b. OFFICE SYMBOL  
(If applicable)  
NA7a. NAME OF MONITORING ORGANIZATION  
Office of Naval Research6c. ADDRESS (City, State, and ZIP Code)  
New England Medical Center Hospital, Inc.  
Box 817, 750 Washington St.  
Boston, MA 021117b. ADDRESS (City, State, and ZIP Code)  
800 N. Quincy Street  
Arlington, VA 22217-50008a. NAME OF FUNDING / SPONSORING  
ORGANIZATION  
Office of Naval Research8b. OFFICE SYMBOL  
(If applicable)  
ONR9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  
N00014-90-5-1560

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street  
Arlington, VA 22217-5000

10. SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NO  
61153NPROJECT  
NO  
RR04108TASK  
NO  
444r003--01WORK UNIT  
ACCESSION NO.11. TITLE (Include Security Classification)  
The development of a mathematical description of the olfactory bulb, validated with intracellular and voltage-sensitive dye recordings: (U)12. PERSONAL AUTHOR(S)  
Kauer, John S.13a. TYPE OF REPORT  
Annual13b. TIME COVERED  
FROM 1/90 TO 1/9114. DATE OF REPORT (Year, Month, Day)  
Jan. 16, 199115. PAGE COUNT  
1516. SUPPLEMENTARY NOTATION  
N/A

17. COSATI CODES

FIELD GROUP SUB-GROUP  
08

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Computational modelling; olfactory system; salamander;  
\* odors \* behavior

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

The objective of this work is to develop a detailed mathematical description of the function of the olfactory epithelium and bulb in the tiger salamander, an animal from which we, and others, have obtained substantial morphological, biochemical, physiological and behavioral information. This project focuses especially on incorporating the information available about the numbers of cells, their connectivities, and the single cell and ensemble physiological responses using voltage-sensitive dye fluorescence, into the model.

## DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited.

20. DISTRIBUTION AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION (U)

22a. NAME OF RESPONSIBLE INDIVIDUAL  
Dr. Ogor Vodyanov

22b. TELEPHONE (Include Area Code) 703-696-4056 OFFICE SYMBOL ONR

PROGRESS REPORT ON CONTRACT N00014-90-J-1560

**CONTRACT TITLE:**

Development of a Mathematical Description of the Olfactory Bulb, Validated with Intracellular and Voltage-Sensitive Dye Recordings.

**RESEARCH OBJECTIVE:**

The objective of this work is to develop a detailed mathematical description of the function of the olfactory epithelium and bulb in the tiger salamander, an animal from which we, and others, have obtained substantial morphological, biochemical, physiological and behavioral information. This project focuses especially on incorporating the information available about the numbers of cells, their connectivities, and the single cell and ensemble physiological responses using voltage-sensitive dye fluorescence, into the model.

**ABSTRACT**

The encoding and neural integration of molecular signals are processes that underly all nervous system functions and are crucial for an organism's ability to recognize chemical compounds in the environment. As a sensitive, broad-spectrum molecular (odor) detector, the sense of smell serves as a model system for studying neuronal, molecular receptors and information processing in synaptic circuits. Central nervous system regions involved in olfaction mediate life-sustaining behaviors such as food-finding and mate recognition in animals and are the sites of devastating pathologies that include epilepsy, schizophrenia, and Alzheimer's disease in humans. Despite the clinical and basic neurobiological importance of this system, little is known about how molecular, odor information is acquired, processed and integrated by the brain.

The long-term aim of this research is to understand how odors are encoded and integrated in the olfactory pathway. To do this, one must study how olfactory neurons function both as single elements and as cooperative members of synaptic networks when stimulated with odors. Our approach has been to develop an animal model (the tiger salamander) which can be electrophysiologically studied after stimulation using controlled odor delivery and from which odor-guided behavior can be obtained. In single cell recording studies this preparation has permitted precise correlation of odors with extra- and intracellular receptor and mitral/tufted unit responses. For recording from cell assemblies and networks we have used the salamander to develop a new method for dynamic recording from the olfactory epithelium and bulb using real-time imaging of voltage-sensitive dye fluorescence.

We have several on-going studies in which we are investigating single cell and ensemble responses from the olfactory pathway. The present report describes the development of a mathematical model of the anatomical properties and physiological events we have observed using the approaches described above. In the course of developing this model, we have found that the process of having to minutely characterize each and every neuronal event in formal mathematical terms has greatly clarified our conceptualization of these processes, specifically highlighting those data which are lacking and emphasizing those events which have been poorly thought through. We think that the ability to develop such a formal description of neurophysiological events in the olfactory bulb in the context of being able to actually record from single cells and the overall network, provides a unique opportunity to validate the mathematical models we, and others, have developed.

**METHODS**

We use a series of linked differential equations which describe the responses in cells making up a somewhat simplified synaptic network representing the olfactory epithelium and bulb of the salamander. The program is written in 'C' and has been run on a variety of computational platforms, including a 16MHz 80386 with a 80387 math coprocessor, a 33 MHz 80386 with a



where  $N_m$  is the number of mitral cells.

Taking into account, for this example, only the change in membrane potential in these two cell types and a membrane decay time constant for each cell, the membrane potential at time  $t$  for granule cell  $i$  coupled to mitral cell  $j$ , will then be:

$$(3) \quad \frac{dg_i}{dt} = -\frac{1}{\tau_g} g_i + \sum_{j=0}^{N_m} a_{ij} m_j$$

and the membrane potential for mitral cell  $i$  coupled to granule cell  $j$  will be:

$$(4) \quad \frac{dm_i}{dt} = -\frac{1}{\tau_m} m_i + \sum_{j=0}^{N_g} b_{ij} g_j$$

where  $\tau_m$  is the membrane decay constant for mitrals

$\tau_g$  is the membrane decay constant for granules

$b_{ij}$  is the granule --> mitral coupling coefficient

$a_{ij}$  is the mitral --> granule coupling coefficient.

Such equations along with equivalent ones representing the connectivities, membrane properties, etc. for all of the other cells are integrated using the method described above and equations for all parameters in all cells are calculated for each time step iteration. We generally calculate sufficient numbers of iterations to represent 2 secs of real time. We have approached these equations as an initial value problem, that is we start all membrane potentials at zero and introduce a spontaneous firing rate in each of the cell types. The model is run using an explicit adaptive step-size algorithm which, as mentioned earlier, adjusts the time step size ( $dt$ ) according to the slope of the function between points; the steeper the slope, the shorter the time step. The model is allowed to first come to a steady state and is then subjected to a simulated stimulus in the form of an electrical or odor-induced depolarization applied to the receptor cells as shown in Figs. 2,3,4 and 6,7.

For reasonable choices of coupling parameters, membrane time constants, firing rates, etc., the model is stable using Euler's integration method, which is, in essence, a linear extrapolation from the present time point to the next one using the slope of the function. We are in the process of evaluating other methods of integration such as, for example, a fourth order Runge-Kutta technique, which examines two points behind and two points ahead of the point of interest, to evaluate differences in efficiency and accuracy among the methods.

## RESULTS

The output of the model is displayed in two formats. Fig. 2 shows displays of the values in each cell of the arrays at various times after the beginning of the run ( $t = xx$ ) in the form of gray scale plots of the matrices. The displays on the left side of the figure represent the values in the receptor cell matrix, with the matrix displayed to schematically represent the actual shape of the ventral receptor sheet in the salamander. The displays on the right side of the figure represent the values in the matrices of the various bulbar components; the periglomerular, glomerular, mitral, and granule elements. The orientation of the bulbar layers in this display approximately conform to their orientation seen real voltage-sensitive dye images. The values in these displays range above and below resting value which is shown by the mid gray value seen uniformly throughout the receptor matrix at  $t = 530$  ms. Gray values lighter than this represent depolarization (white represents spikes), darker grays represent hyperpolarization.

In this trial a simulated electrical stimulus was applied to the locations shown on the

receptor sheet at  $t = 500$  ms (first row, white squares). As shown at  $t = 530$  ms, this stimulus elicited depolarization of periglomerular cells (with spikes) and of glomerular elements. At  $t = 546$  ms depolarization appeared in mitral and granule cells and at  $t = 580$ , the granule cells showed widespread depolarization and the mitrals hyperpolarized.

The second display format is shown in Figs. 3,4 and 6,7. These figures show the changes in membrane potential for particular cells as if one were recording from them using an intracellular electrode. Recordings from one receptor cell (of 1000), one periglomerular cell (of 10), one glomerular element (of 20), and one granule cell (of 1000) are shown at the tops of these figures. Simulated recordings from all ten mitrals are shown toward the bottoms of the figures.

In Figs. 3 and 4 a simulated electrical stimulation was applied to receptor cells 300-700 at low and high intensities. With a low intensity, two spikes are elicited from the pg cell, and a single spike is elicited in mitral cells 2-6 followed by a hyperpolarization. Fig. 4 shows that with high intensity stimulation more spikes are elicited from pg's and mitral cells, a spike now appears in the granule cell, and the mitral hyperpolarization is deeper and longer lasting. These simulated responses are to be compared with real intracellular recordings from salamander mitral/tufted cells shown in Fig. 5 (made by K.A. Hamilton in this lab). Notice that in these records, with more intense stimulation, more spikes and a deeper and longer lasting hyperpolarization are elicited.

In Figs. 6 and 7 a simulated, heterogeneously distributed odor stimulation at high and low intensities is applied to the receptor matrix (see display of receptor matrix at bottom left of each fig). Here the responses in the mitral cells are more complex than with electrical stimulation, consisting of brief bursts of spikes, brief, slight hyperpolarizations, long lasting depolarizations, and longer-lasting following hyperpolarizations. With higher intensity odor stimulation there are actually fewer spikes elicited from mitral cells than with the lower intensity. These responses are surprisingly similar to those observed in real intracellular recordings from mitral/tufted cells at different odor intensities as shown in Fig. 8 (made by K.A. Hamilton in this lab) and are consistent with the concept of concentration tuning we had previously developed on the basis of extracellular studies.

We have been greatly encouraged by these results because we initially put into the model parameters that made the most physiological sense in order to get results similar to those we had obtained from the real system under conditions of **electrical stimulation**. Once the model functioned with these parameters we then tested it with simulated odor stimuli and obtained the results shown here **without manipulating the parameters**. That is the model was not setup using knowledge about the odor responses, but only using knowledge about the electrical responses; only then was it tested for its odor responsivity. As one can see by comparing Figs. 6 and 7 with Fig. 8, the similarity is striking.

We now think we have a reasonably good starting model with which to examine the effects of manipulating a wide variety of physiological parameters such as the effects of changing synaptic efficacy using pharmacological agents.

#### **PUBLICATIONS AND REPORTS (Year 1):**

Kauer, J.S., Neff, S.R., Hamilton, K.A., Cinelli, A.R. (1991) The Salamander Olfactory Pathway: Visualizing and Modelling Circuit Activity. In: Davis, J. and Eichenbaum, H. (eds.), Olfaction as a Model System for Computational Neuroscience. MIT Press. (in press).

White, J., Neff, S.N. Cinelli, A.R., Kauer, J.S. (1990) Modelling the salamander olfactory bulb: single cell and network interactions. Neurosci. Abst., abstract.

White, J., Neff, S.N., Hamilton, K.A., Kauer, J.S. (1991) A comprehensive model of the peripheral olfactory pathway of the salamander. J. Biophys. (in preparation)








FIG. 1.



t = 502 msec





t = 530 msec



t = 534 msec



t = 546 msec



t = 580 msec





FIG. 2

# Low Electrical Stimulus Intensity




FIG. 3



# High Electrical Stimulus Intensity




FIG. 4





FIG. 5


# Low Odor Concentration



Reproduced From  
Best Available Copy

FIG. 6

# High Odor Concentration



Reproduced From  
Best Available Copy

FIG. 7




FIG. 8

OLFACTORY DISCRIMINATION PROGRAM

- ✓ Dr. Barry W. Ache  
Whitney Laboratory  
University of Florida  
9505 Ocean Shore Blvd.  
St. Augustine, FL 32086-8623
- ✓ Dr. Richard C. Bruch  
Dept. of Neurobiology & Physiology  
Northwestern University  
O. T. Hegan Hall  
Evanston, IL 60208
- ✓ Dr. John Caprio  
Dept. of Zoology & Physiology  
Louisiana State University  
Baton Rouge, Louisiana 70803-1725
- ✓ Dr. Vincent Dionne  
Dept. of Pharmacology  
M-036, School of Medicine  
University of California  
La Jolla, CA 92093
- ✓ Dr. Stuart Firestein  
Section of Neuroanatomy  
Yale University  
School of Medicine  
C303 Sterling Hall of Medicine  
333 Cedar Street  
New Haven, Connecticut 06510
- ✓ Dr. Stephen P. Fracek, Jr.  
Dept. of Biological Sciences  
University of North Texas  
P.O. Box 5218  
Denton, TX 76203-5218
- ✓ Dr. William F. Gilly  
Hopkins Marine Station  
Stanford University  
Pacific Grove, CA 93950
- Dr. John S. Kauer  
Dept. of Neurosurgery  
New England Medical Center  
750 Washington Street  
Boston, MA 02111
- ✓ Dr. Mimi A. R. Koehl  
University of California  
540 Banway Building  
Berkeley, CA 94720
- ✓ Dr. Michael Leon  
Dept. of Psychobiology  
University of California  
19172 Jamboree  
Irvine, CA 92717
- ✓ Dr. Donal Manahan  
Dept. of Biology  
Univ. of So. California  
University Park  
Los Angeles, CA 90089
- ✓ Dr. Thomas H. Morton  
Department of Chemistry  
University of California  
Riverside, CA 92521-0217
- ✓ Dr. W. Charles Neely  
Department of Chemistry  
Auburn University  
Auburn, Alabama 36849-3501
- ✓ Dr. Dennis E. Rhoads  
Dept. of Biochemistry  
and Biophysics  
University of Rhode Island  
70 Lower College Road  
Kingston, RI 02881
- ✓ Dr. Jeanette Yen  
Marine Science Research  
Center  
SUNY  
Stony Brook, NY 10019

Olfactory Discrimination Program

Final and Technical Reports

ADMINISTRATORS

- ✓ Dr. Igor Vodyanoy, Code 1141SB (2 copies)  
Scientific Officer, Biophysics  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000
- ✓ Administrator (2 copies) (Enclose DTIC Form 50)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314
- ✓ Administrative Contracting Officer  
ONR Resident Representative  
(address varies - obtain from contract or your business office.)
- ✓ Dr. Randall Alberte, Code 1141MB  
Scientific Officer  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000
- ✓ Dr. Joel Davis, Code 1142BI  
Program Manager, Biological Intelligence  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000
- ✓ Program Manager, Code 125  
Biological/Human Factors Division  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000
- ✓ Program Manager Defense Technical Support  
Technology Directorate  
Office of Naval Technology, Code 223  
800 N. Quincy Street  
Arlington, VA 22217-5000

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

- ✓ Commander  
Chemical and Biological Sciences Division

Research Army Research Office, P.O. Box 1221  
Research Triangle Park, NC 27709

Olfactory Discrimination Program

DoD ACTIVITIES (continued)

- ✓ Head, Biomolecular Engineering Branch  
Code 6190  
Naval Research Laboratory  
Washington, DC 20375
- ✓ Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332
- ✓ Dr. Tove  
4300 S. Miami Boulevard  
Research Triangle Park, NC 27709

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)  
Attn: Technical Information Division, Code 2627  
Washington, DC 20375